

## References for LDRD DR Proposal 20060049DR

- [1] S. S. Adler et al. [PHENIX Collaboration], "Elliptic flow of identified hadrons in Au + Au collisions at  $\sqrt{s_{NN}} = 200$  GeV", Phys. Rev. Lett. **91**, 182301 (2003).
- [2] S. S. Adler et al. [PHENIX Collaboration], "Suppressed  $\pi^0$  production at large transverse momentum in central Au + Au collisions at  $\sqrt{s_{NN}} = 200$  GeV", Phys. Rev. Lett. **91**, 072301 (2003).
- [3] I. Vitev and M. Gyulassy, "High-p<sub>T</sub> tomography of d + Au and Au + Au at SPS, RHIC, and LHC", Phys. Rev. Lett. **89**, 252301 (2002).
- [4] N. Brambilla et al., "Heavy Quarkonium Physics," e-Print hep-ph/0412158 (2004).
- [5] F. Cooper, E. Mottola and G. C. Nayak, "Minijet initial conditions for non-equilibrium parton evolution at RHIC and LHC", Phys. Lett. B **555**, 181 (2003).
- [6] M. Gyulassy, P. Levai and I. Vitev, "Non-Abelian energy loss at finite opacity", Phys. Rev. Lett. **85**, 5535 (2000).
- [7] M. Djordjevic and M. Gyulassy, "Heavy quark radiative energy loss in QCD matter", Nucl. Phys. A **733**, 265 (2004).
- [8] F. Karsch, "Deconfinement and Quarkonium Suppression," e-Print hep-lat/0502014 (2005).
- [9] P. Petreczky, "QCD Thermodynamics on Lattice," e-Print hep-lat/0409139 (2004).
- [10] S. S. Adler et al. [PHENIX Collaboration], "Measurement of single electron event anisotropy in Au+Au Collisions at  $\sqrt{s_{NN}} = 200$  GeV," e-Print: nucl-ex/0502009 (2005).
- [11] T. Tabaru [PHENIX Collaboration], "Heavy quark and  $J/\psi$  production at RHIC / PHENIX", International Conference on Physics and Astrophysics of Quark Gluon Plasma, Kolkata, India, February 8-12, 2005.
- [12] D.M. Jansen et al., "Measurement of the Bottom-Quark Production Cross Section in 800 GeV/c Proton-Gold Collisions", Phys. Rev. Lett. **74**, 3118 (1995).
- [13] P.L. McGaughey et al., "Study of Open Charm Production at RHIC," LDRD ER Project #20020043ER, completed in 2004.